



Influence of different milking methods on milk quality based on somatic cell count and basic composition

Jana Zahumenská^{2,D}, František Zigo^{1,F}, Mariana Kováčová^{2,C}, Silvia Ondrašovičová^{3,E}, Vladimír Hisira^{4,B}, Tomáš Mihok^{1,B}, Jana Výrostková^{2,E}, Zuzana Farkašová^{1,B}

¹ Department of Animal Nutrition and Husbandry, University of Veterinary Medicine and Pharmacy, Košice, Slovak Republic

² Department of Hygiene, Technology and Health Food Safety, University of Veterinary Medicine and Pharmacy, Košice, Slovak Republic

³ Department of Biology and Physiology, University of Veterinary Medicine and Pharmacy, Košice, Slovak Republic

⁴ University Veterinary Hospital, Clinic of Ruminants, University of Veterinary Medicine and Pharmacy, Košice, Slovak Republic

A – Research concept and design, B – Collection and/or assembly of data, C – Data analysis and interpretation,

D – Writing the article, E – Critical revision of the article, F – Final approval of the article

Zahumenská J, Zigo F, Kováčová M, Ondrašovičová S, Hisira V, Mihok T, Výrostková J, Farkašová Z. The influence of different milking methods on milk quality based on somatic cell count and basic composition. *Ann Agric Environ Med.* doi: 10.26444/aaem/187170

Abstract

Introduction and Objective. Correlations between the number of milk somatic cells (SCC), the number of microorganisms, and the content of basic components of milk were studied on five farms (F1–F5) with cows of the same breed, but with different milking systems.

Materials and Method. From each farm, 50 Holstein Friesian milk samples were collected once a month (250 samples/month; n=3,000) during March 2022 – February 2023. Samples from farms F1 and F5 were tested for fat, protein, lactose, no fat dry matter content (FTIR spectroscopy), for the SCC (Fossomatic 7), and for the differential cells (Vetscan DC-Q).

Results. The highest fat content was confirmed on farm F5 (3.85 ± 1.70%) and F4 (3.82 ± 0.21%) with automatic milking system (AMS). However, from the point of view of protein content, these farms showed slightly lower values (<0.05). F1 did not meet the minimum required amount for fat content (2.84 ± 0.81%) set by the legislation of the Slovakia. The comparison shows that there is not much difference in cell size between healthy cells and mastitis cells. The average size of healthy cells was approximately 8.77 ± 0.49 µm. In the monitored period, the average values determined were at the level of 292,000/mL (5.46 ± 0.72 log₁₀ SCC) in cow milk samples, while for the rest of the year, the values remained at 256,000/mL (5.40 ± 0.80 log₁₀ SCC). F1 was categorized as a positive farm with a high TLC (total milk leucocyte count) concentration (5.58 log₁₀ cells/mL, 406.65 ± 53.80 × 10³ cells/mL) and a predominant NEU fraction (61%). Farms F2, F4, and F5 were classified as negative farms (TLC was 4.70 ± 0.26 log₁₀ cells/mL).

Conclusions. According to the results, the size of SCCs in healthy milk does not differ from SCCs found in mastitis milk. From the results, it can be concluded that the transition to the latest generation of robotic milking method can positively affect milk production and its quality.

Key words

somatic cells, size, milk components, method of milking, seasonality, quality, mastitis

INTRODUCTION

Milk contains several types and different numbers of cells which are a very important criterion of udder health, and an indicator of the technological, hygienic and health quality of milk [1]. Cells enter the milk either from the udder itself or from the bloodstream of the dairy cow.

Somatic cells include mainly milk-secreting epithelial cells that have been shed from the lining of the gland, and white blood cells (leukocytes – monocytes, granulocytes, and lymphocytes) that have entered the mammary gland in response to injury or infection [2].

Somatic cell count (SCC) in milk is highly correlated with udder health, and somatic cells are generally the standard in its diagnosis [3, 4].

Some authors [5] state that mastitis is an inflammatory disease of the mammary gland, which is characterized by physical, chemical and microbial changes, as well as an increase in the number of somatic cells and other changes in the composition of milk. Mastitis is among the most economically serious livestock diseases. Due to its high prevalence and negative impact on the economy, mastitis is a major problem in the dairy industry [6]. This problem is widespread and causes a decrease in the quantity as well as the quality of milk produced. It is estimated that cows affected by subclinical mastitis produce 25–42% less milk than healthy cows [7, 8, 9].

The SCC increases as leukocytes – particularly polymorphonuclear leukocytes (PMNs) are recruited from the blood into the mammary gland to phagocytize invading organisms. This immune response is crucial for preventing further infection by pathogens. Because the magnitude of the SCC is closely linked to the extent of inflammation in the mammary gland, cows with the lowest averages of SCC during lactation are deemed to have the highest resistance to intramammary infection (IMI) and mastitis [10].

✉ Address for correspondence: František Zigo, Department of Animal Nutrition and Husbandry, University of Veterinary Medicine and Pharmacy, Komenského 73, 041 81 Košice, Slovak Republic
E-mail: frantisek.zigo@uvlf.sk

In 2017, a new tool – the differential somatic cell count (DSCC) – was described for determining the proportion of PMNs and lymphocytes in somatic cells during monthly DHI testing (Dairy Herd Information). Given that lymphocytes typically constitute a small proportion of milk cells, fluctuations in DSCC primarily reflect changes in PMN levels [11].

In cows affected by mastitis, almost all the basic components of milk are changed. Therefore, data on the composition of milk are often used for general estimation of animal health and udder condition, as well as an indicator for detecting subclinical mastitis, feeding errors, causes of metabolic disorders. The composition of milk is influenced by other factors, such as breed, number of lactations, stage of lactation, technology of housing, season, feeding, environmental factors [12].

However, few studies have evaluated the association between breed and IMI, or mastitis associated pathogens. Previous studies found that some breed variation may exist, for example, *Staphylococcus haemolyticus* is more common in quarter milk samples from Swedish Holsteins compared with Swedish Reds [13], and *Staphylococcus devriesei* was more common on the teat ends of red and white Holstein Friesian cows compared with black and white Holstein Friesian cows [14]. Conversely, another study identified no impact of breed when evaluating pathogens associated with clinical mastitis among Swedish dairy cows [15, 16].

Another influence on milk quality is the method of milking (manual, machine). In general, automatic milking system (AMS) has been adopted as a realistic alternative for milking in the 'traditional' milking parlour. Systems have gradually been improved and, maybe even more importantly, farmers have become more familiar with their potential and limitations, both technically and in herd management. The number of farms milking with AMS has increased worldwide [17]. The technical design and the functionality of an AMS could influence the behaviour and milking characteristics of dairy cows, both directly and indirectly.

All developed countries use the milk SCC as a marker to monitor the prevalence of mastitis in dairy herds. Among all the different screening tests for milk quality, it is the number of somatic cells in milk that is the most effective method for detecting the subclinical form of mastitis [18]. Regulation of the European Parliament and Council No. 853/2004 of 29 April 2004, established special hygiene regulations for food of animal origin, i.e. the criterion for the number of somatic cells in raw cow's milk in a pool sample is a maximum of 400,000 cells/mL [19]. The number of somatic cells is a mandatory indicator in Europe. In the USA, the legal maximum SCC count for Grade A farm bulk milk is 750,000 cells/mL; this limit is high compared to many international standards. Much of Europe, Australia and New Zealand, has a limit of 400,000 cells/mL, and Canada a limit of 500,000 cells/mL of raw milk. Milk SCC is a diagnostic figure for subclinical mastitis [20].

Slovakia is a landlocked central European country with a climate that can be described as a typical European continental [21].

Cow's milk production is an important and traditional sector of agricultural primary production in Slovakia. The current article presents results that provide a framework picture of the hygienic quality of raw cow's milk on five monitored Slovak farms for a period of one year (March 2002 – February 2023). The aim of the study, therefore, was to investigate the quality of milk and to compare the

influence of the number of somatic cells on the content of milk components on selected farms. The location, effect of the season and method of milking on the mentioned parameters were taken into account. Somatic cells were analyzed in terms of morphological characteristics – size and number of cells, i.e. total number of milk leukocytes (TLC), such as Lymphocytes, Macrophages, Neutrophils.

MATERIALS AND METHOD

Individual samples of raw cow's milk were obtained from five farms (F1 – F5) from different regions of Slovakia (Fig. 1). Dairy cows are stabled on all farms with deep bedding, the food regime was the same on all farms.

Sampling was based on the real conditions of breeding practice in the Slovak Republic. Individual milk samples ($n = 3,000$) were collected from Holstein Friesian cattle. Fifty samples of milk of the Holstein Friesian cattle were taken from each farm once a month (250 samples/month) for a period of one year (March 2022 – February 2023). Dairy cows were not treated with antibiotics for mastitis or any other disease. Dairy cows had first to fifth lactation (2.67 ± 1.52) and the 12-month average DIM on farms was 270–300 days.

Farm F1 (500 dairy cows) is located in the southeast of the Slovak Republic, has a lowland-hilly character (200 m a.s.l.). The territory belongs to the northern temperate zone with average annual temperatures of around 10°C. Southwest of the city of Košice (the metropolis of eastern Slovakia), F2 is located (400 dairy cows) which also belongs to the Košice region. F3 (300 dairy cows) is located in the southeastern part of the Republic (170 m a.s.l.) and borders the Republic of Hungary. All three farms use Boumatic parallel milking equipment. F4 and F5 (350 dairy cows) were among the first unified farming cooperatives in Slovakia, located in the Banskobystrický Region at the foot of the Poľana mountain range at an altitude of 400 m a.s.l. The farms use a robotic method of milking (F4 – Lely Astronaut A4; F5 – Lely Astronaut A4).



Figure 1. Geographic location of sampling from five farms (F1 – F5)

At each farm, feeding was carried out using complete mixtures of feed rations, the so-called TMR (total mixed ration). One of the biggest advantages of TMR is the stable composition of the ration.

The milk samples were transported to the laboratory in a refrigerator (4°C) and processed for analysis within 12 hours. Samples from farms F1 – F5 were delivered to the Examinálna testing laboratory (Žilina, Slovak Republic), which is the central laboratory for testing raw milk in Slovakia.

Determining the basic components of milk. In samples from all farms fat, protein, but not fat solids (SNF) content, were tested by FTIR spectroscopy using MilkoScan FT 6000 (Foss Analytical A/S, Hillerød, Denmark), according to the equipment manual.

Somatic Cells Analysis. SCC in samples from all farms was analyzed with the use of a Fossomatic 7 flow cytometer (FOSS, Denmark) according to ISO 13366-2 [22]. All examinations are accredited by SL Examinála by the Slovak National Accreditation Service according to ISO/IEC 17025.

Differential cells were also counted using a Vetscan DC-Q milk analyzer (AAD Advanced Animal Diagnostics, NC, USA) that provides the concentration (cells/mL) of total milk leucocyte count (TLC), defined as SCC without epithelial cells. The results of SCC were transformed to logarithmic form. Differential cells were used to calculate the differential somatic cell count (DSCC), defined as the sum of neutrophils (NEU) and lymphocytes (LYM) as a percentage of total SCC. According to many authors [23, 24], this index (DSCC) increases during intramammary infections. The Vetscan DC-Q milk analyzer uses fluorescence imaging, as described by Godden [25], and interprets the results through secret algorithms that identify intramammary infections.

Microbial analyses. Preparation of milk samples, initial suspension, and decimal dilutions were prepared according to ISO 6887-5 (2011) [26].

All procedures were performed according to Lobacz et al. [27]. The milk samples were decimally diluted in peptone physiological salt solution, and an inoculum of 100 µL was used. Colonies on plate count agar (Oxoid Ltd., Basingstoke, UK) were counted after 72 h incubation at 30 °C.

Subsequently, the Pure Milk Test (PM test) was used – for farm diagnosis of mastitis (LabMediaServis, CZ). The PM

test, which is used for farm diagnosis of mastitis, is a quick and very effective solution for correct diagnosis and, thanks to this, targeted treatment of the causative agents of the mammary glands in dairy cows. The diagnostic set included a three-sector petri dish with special chromogenic agars. The milk sample was applied using an inoculation stick to the surface of all three agar Petri dishes, and incubated for 22–26 hours at 37.5 °C. Subsequently, the type of causative agent of mastitis was determined using the Atlas of Agents [28]. The results were expressed in several colony forming units (CFU/mL).

Statistical analysis. In the statistical evaluation, the differences were compared in milk quality between the location, influence of the season, and the method of milking. The results were evaluated using MS Excel (Microsoft, Redmond, Washington, USA). Basic statistical characteristics, such as mean, and standard deviation, were calculated for numerical data.

Statistical analysis was performed by one-way analysis of variance. ANOVA and Tukey's test for multiple comparisons of means with a confidence interval set at 95% were performed using GraphPad Prism statistical software 8.3.0.538 (GraphPad Software, San Diego, CA, USA).

RESULTS

Correlations between milk somatic cell count (SCC), the number of microorganisms, and the content of the basic components of milk were studied on five farms with cows of the same breed in different seasons, and locations, and the method of milking. Annual data (year 2022/2023) were analysed which meant 3,000 milk samples.

The average content of fat, protein, and solids not fat is shown in Table 1.

Table 1. Mean ± and their standard deviation (SD) of the basic indicators of cow's milk, depending on the month of milking

Season	Spring			Summer			Autumn			Winter		Avg. per year		
	March	Apr	May	Jun	July	Aug	Sept	Oct	Nov	Dec	Jan		Feb	
FAT (%)	F1	2.90	2.90	3.10	3.00	3.00	2.71	2.21	2.35	2.80	2.80	3.10	3.30	2.84 ± 0.81 ^c
	F2	3.20	3.34	3.43	3.41	3.32	3.46	3.37	3.00	3.62	3.70	3.11	3.30	3.35 ± 0.18 ^{b,c}
	F3	3.30	3.22	3.30	3.39	3.25	3.32	3.60	3.50	2.95	2.80	2.9	2.98	3.20 ± 0.23 ^{b,c}
	F4	4.03	3.90	3.80	3.67	3.95	4.12	4.10	3.70	3.54	3.40	3.80	3.89	3.82 ± 0.21 ^{a,b}
	F5	4.03	3.98	3.80	3.85	3.80	3.95	4.00	9.96	3.20	3.50	4.10	4.05	3.85 ± 1.70 ^a
														<0.001
PRO (%)	F1	3.60	3.48	3.62	3.59	3.44	3.68	3.55	3.49	3.47	3.52	3.47	3.56	3.53 ± 0.07 ^a
	F2	3.28	3.36	3.33	3.31	3.56	3.55	3.52	3.47	3.27	3.47	3.45	3.52	3.42 ± 0.10 ^{c,a}
	F3	3.37	3.25	3.21	3.37	3.38	3.29	3.00	3.37	3.53	3.55	3.6	3.52	3.37 ± 0.16 ^{b,c}
	F4	3.40	3.40	3.30	3.20	3.30	3.40	3.45	3.30	3.35	3.30	3.40	3.35	3.34 ± 0.06 ^{b,c}
	F5	3.35	3.37	3.30	3.18	3.20	3.30	3.30	3.22	3.10	3.28	3.30	3.34	3.27 ± 0.07 ^b
														<0.001
SNF (%)	F1	9.80	9.50	9.84	9.81	9.41	10.70	9.68	9.53	6.44	9.66	9.45	9.71	9.46 ± 0.96 ^a
	F2	9.12	9.10	9.17	9.11	9.50	9.71	9.59	9.48	8.92	9.47	9.40	9.49	9.33 ± 0.23 ^a
	F3	9.12	8.85	8.75	9.19	9.21	8.96	8.20	9.60	9.64	9.55	8.36	9.61	9.09 ± 0.46 ^a
	F4	9.18	9.20	9.06	8.94	9.13	9.20	9.10	9.00	8.70	8.62	8.89	8.55	8.96 ± 0.21 ^b
	F5	9.17	9.20	9.16	9.00	9.15	9.11	9.19	8.97	9.19	9.16	9.19	9.20	9.14 ± 0.07 ^a
														<0.001

PRO – protein; SNF – solids not fat

a, b, c – within a row, means without a common superscript differ ($P < 0.05$)

The month of milking significantly affected all quality traits, as above. The content of fat, and not fat solids, increased with the time of lactation ($P < 0.05$).

The annual average showed statistically significant ($P < 0.05$) positive correlation between the content of fat, protein and solids not fat content (Figure 2).

The highest and almost identical fat content was confirmed on farm F5 ($3.85 \pm 1.70\%$) and F4 ($3.82 \pm 0.21\%$) which used the Lely Astronaut robot method of milking. However, from the point of view of protein content, these farms showed slightly lower values (< 0.05).

F1 did not meet the minimum required amount for fat content ($2.84 \pm 0.81\%$) set by the legislation of the Slovak Republic.

Within the seasons, the values for fat were highest in spring and summer ($3.48 \pm 0.44\%$), proteins ($3.38 \pm 0.14\%$) and SNF (9.34 ± 0.36) only in the summer months.




Figure 2. Dependence between fat content, fat-free solids and protein content in milk on selected farms during the monitored period

In this study, statistical significance was demonstrated between SCC and their size ($P < 0.005$), as well as between SCC and fat content in all farms ($P < 0.005$). The relationship between cell size and seasons (Fig. 3) was not statistically proven ($P > 0.05$). Changes in the number of somatic cells within the seasons are presented in Figure 3.




Figure 3. Average number of somatic cells on five farms during the monitored period

The comparison shows that there is little difference in cell size between cells from healthy and infected mammary glands. The average size of healthy cows ranged from approximately $8.77 \pm 0.49 \mu\text{m}$.

The smallest size of SCC ($6 \mu\text{m}$) was shown by the sample from F5 in the spring, the highest size of SCC ($12.13 \mu\text{m}$) was recorded on F2, also in spring (Tab. 2). This means that the size of somatic cells is not decisive for health.

The correlation between SCC and cell size on monitored farms is shown in Figure 4.




Figure 4. Correlation between the number of somatic cells and their size in milk on selected farms during the monitored period

Using an analyser, it can be assessed that in the negative sample there were mainly epithelial cells of the milk ducts, and a small amount of leukocytes. Table 3 shows the results of analyses using the Vetscan DC-Q milk analyzer.

The F1 farm was categorized as a positive farm with a high TLC concentration ($5.58 \log_{10} \text{ cells/mL}$, $406.65 \pm 53.80 \times 10^3 \text{ cells/mL}$) and a predominant NEU fraction (61%). The results from F3 reached borderline values ($386.17 \pm 21.36 \times 10^3 \text{ cells/mL}$) with a TLC of $4.9 \log_{10} \text{ cells/mL}$ and a higher NEU percentage of 63.10%. Farms F2, F4, and F5 were classified as negative farms. TLC, on average, was $4.70 \pm 0.26 \log_{10} \text{ cells/mL}$, while the percentage of NEU ($52.0 \pm 14.15\%$) was lower than the percentage of positive cows. In addition, the number of MAC in the milk of F2, F4, and F5 cows was higher (19%) than that of F1 milk (8.70%).

The aggregated microbiological quality of raw cow's milk samples from five farms is been presented in Figure 5. The results are satisfactory, although on F1 ($90.16 \pm 8.14 \times 10^3 \text{ CFU/mL}$; $4.94 \pm 0.03 \log \text{ CFU/mL}$) bordering on the upper limit for maximum limits (100,000 CFU/mL), which can be justified mainly by the current state of hygiene in the process of obtaining milk. The results from the F1 and F2 farms show the occurrence of mastitis in dairy cows.

F4 was the opposite, where the results of the arithmetic mean of the tested samples were $26.33 \pm 18.19 \times 10^3 \text{ CFU/mL}$ ($4.33 \pm 0.25 \log \text{ CFU/mL}$) because very strict hygienic criteria for milking and housing dairy cows are observed on the farm.




Figure 5. Total bacteria count (log CFU/mL) in samples of raw milk

The results show the lowest number of total bacteria count (TBC) on farms with AMS ($4.45 \pm 0.19 \log \text{ CFU/mL}$) (Fig. 6). In May – July, values were stable on farms, the highest increase was during the transition from winter to spring. Whereas in general the highest incidence of diseases were recorded in August and September, in the current study there was an increase in autumn.

The study demonstrates a statistical significance between SCC and TBC within the monitored period on the monitored farms ($P < 0.005$).

Table 2. Relationship between SCC and cell size on five farms with different milking methods during the year

Farm	F1		F2		F3		F4		F5	
Value	SCC	Size	SCC	Size	SCC	Size	SCC	Size	SCC	Size
MIN	4.96	6.34	4.92	6.5	4.58	6.63	4.07	6.12	3.77	6.12
MAX	6.53	10.98	6.70	12.13	6.73	10.72	5.47	11.05	5.47	11.05
Mean±SD	5.60 ± 0.86		5.40 ± 0.74		5.58 ± 0.80		5.27 ± 0.65		5.26 ± 0.61	
AVG	406.65 ± 53.80	8.66 ± 2.32	256.32 ± 59.70	9.31 ± 2.81	386.17 ± 21.36	9.38 ± 0.84	187.16 ± 32.20	8.25 ± 2.75	183.83 ± 21.83	8.25 ± 2.75
pValue	<0.05		<0.05		<0.05		<0.05		<0.05	

SCC – somatic cell count (x 10³ cells/mL); Size – average cell size (µm); Size – average cell size (µm)


AVG – average (x 10³ cells/mL)

Mean ± SD – means and their standard deviations of log₁₀ SCC in cow milk samples on different months of milking

Table 3. Results of the total milk leucocyte count and their differential cell count (Lymphocytes, Macrophages, Neutrophils) in milk on various farms in Slovakia

Value	TLC [log ₁₀ cells/mL]	Lymphocytes [%]	Macrophages [%]	Neutrophils [%]
AVG	4.70 ± 0.26	26 ± 14.90	23.70 ± 15.59	52.0 ± 14.15
Min	4.60	1.0	1.0	1.0
Max	5.58	68.20	63.90	90.50

TLC – total milk leucocyte count; AVG – average (log₁₀ cells/mL)

**Figure 6.** Correlation between the number of somatic cells and total bacteria count

DISCUSSION

The importance of the influence of the season on the quality of milk has been discussed and confirmed by various authors [12] because geographical and climatic conditions cannot be influenced. The time of year is often associated with different eating regimes, which allow for sufficient intake, good digestion for metabolism; on the other hand, they can affect the composition of milk. Some authors report that changes in milk composition are more related to feeding rather than genetic factors; hence, for better correlations among variables (composition) the food intake is more important than the content of nutritive matters in a diet. Nevertheless, the season of the year considerably affects the food intake [29].

SCC is influenced by many factors, such as cattle breed, level of milk production, stage and order of lactation, season, month of measurement, individual and environmental factors, teat and udder morphology, as well as management practices [30, 31, 32, 33].

The results obtained in the current study confirm [34] that from the turn of April – May to the turn of September – October, a slight increase in the number of somatic cells are observed. When dairy cows are housed on deep bedding, the increase is also noticeable during the winter months. This fact has also been confirmed by other authors [35],

who in their study for 2015, demonstrate an increase in the number of somatic cells between May – August to a value approaching 280,000 cells/mL, while for the rest of the year the values remained at 230,000 cells/mL. In the monitored period from May – August in the current study, the average values determined were at the level of 292,000/mL (5.46 ± 0.72 log₁₀ SCC) in cow milk samples, while for the rest of the year the values remained at 256,000/mL (5.40 ± 0.80 log₁₀ SCC). Other authors [36] state that somatic cells reach their highest values in August and September. However, the main factor affecting PSB in milk is infection of the mammary gland [37].

An important factor in describing the efficiency of an AMS is the cow-individual SCC, is that it is not influenced mainly by the milking technique [38], but that the changeover of the milking system could have an effect on the SCC of a cow. In one study [39], the authors found that the highest SCC occurred in the first six months after the introduction of an AMS, after which the SCC normalized and stabilized. Another study [40] compared two AMS from different manufacturers regarding milk yield, milking frequency, milking interval, teat-cup attachment success rate, and length of milking procedure. They found differences in teat-cup attachment success, duration of several milking phases, and milking frequency regarding different AMS.

Therefore, one of the objectives of the current study was to discover whether a different milking method or the latest generation AMS has advantages in terms of efficiency and animal health, compared to the previous model of the same manufacturer.

One research [41] revealed that high SCS levels do not necessarily negatively affect milk production as long as DSCC is also high [42]. This finding suggests that cows with increased SCS and DSCC may be in the early stages of infection when the infection is well controlled by abundant PMNs. Conversely, when SCS is high while DSCC is low, indicates chronic infection and cow productivity is significantly impaired.

The total bacteria count (TBC) is considered an indicator of the microbiological purity of milk and thus of the environment [43]. Concentrations of colony-forming units can be expressed using logarithmic notation, where the value shown is the base 10 logarithm of the concentration.

There are many management factors that play the most important role in the development of infectious diseases like mastitis in dairy animals, among them unsanitary conditions are more important in increasing the chances of intramammary infection (IMI) and leading to high SCC [44].

The study also evaluated the prevalence of microbial organisms and the association with the number of milk

somatic cells and the persistence of infection [16]. Through the PM test, the presence of colonies of pathogens causing mastitis was diagnosed. Subsequently, the type of causative agent of mastitis was determined using the Atlas of Agents (*Pseudomonas aeruginosa*, *Klebsiella oxytoca*, *Streptococcus dysgalactiae*, *Enterococcus faecalis*, *Streptococcus uberis*, *Staphylococcus chromogenes*, *Staphylococcus spp.*). However, these results would require more detailed research on the prevalence of pathogens and their elimination in breeding.

CONCLUSIONS

Based on the results obtained, it can be concluded that the hygienic quality of the raw cow's milk produced on the selected Slovak farms is at a relatively very good level, with the majority of samples (85%) meeting all legislative limits.

The results confirmed a higher occurrence of microorganisms in milk in the summer season on two of the farms, which were attributed to climatic conditions, unsuitable litter, insufficient animal hygiene, as well as inappropriate feed rations from a qualitative and quantitative point of view. The insufficient nutrition resulted in an increase in the number of somatic cells in milk.

The results of this study confirmed that the production and quality of milk are significantly influenced by the already mentioned external and internal factors, and also confirmed that the size of somatic cells in healthy udder does not differ from the somatic cells found in an infected udder. Furthermore, it was shown that the size of the somatic cells was not decisive in the occurrence of mastitis, as the differences in cell sizes were small.

From the results, it can be concluded that the transition to the latest generation robotic method of milking can positively affect milk production and its quality. The results of this study indicate a higher level of milk quality influenced by milking time, milk flow and milking interval. Even if the milking time is reduced, the milk yield, milk flow and milking interval will increase in the robot. Significant differences were found in milk quality between the two systems.

For the management of the health status of dairy cows and the further processing of milk, further investigation certainly makes sense, but it requires a deeper investigation in relation to the technological quality of raw milk, as well as from the point of view of the possible presence of microorganisms involved in the development of inflammation of the udders of cows.

Acknowledgements

Slovak Research and Development Agency under Contract no. APVV-22-0457 and by the Research and Development Agency under Contract No. SK-PL-230066.

REFERENCES

1. Frančáková H, Čuboň J, Michalčová, A. Evaluation of agricultural products. SUA Nitra, 2005;178. ISBN 80-8069-471-0.
2. Sharma N, Singh NK, Bhadwal MS. Relationship of somatic cell count and mastitis: An overview. Asian-Australasian J Animal Sci. 2011;24(3):429–438. <https://doi.org/10.5713/ajas.2011.10233>
3. Pyörälä S. Indicators of inflammation in the diagnosis of mastitis. In Vet Res. 2003;34:565–578. <https://doi.org/10.1051/vetres:2003026>
4. Vršková M, Tančin V, Uhrinčat M, et al. Survey of somatic cell counts and milk composition in bulk milk of ewes in dairy practice. Slovak J Food Sci. 2021;15:151, ISSN 1338–9971. <https://doi.org/10.5219/1325>
5. Drahošová K, Drončovský M. Mastitis – the most common cause of reduced milk quality. In Mliekarstvo, 2004;35(4):7.
6. Hogeveen H, Steeneveld W, Wolf CA. Production diseases reduce the efficiency of dairy production: A review of the results, methods, and approaches regarding the economics of mastitis. Ann Rev Res Econom. 2019;11:289–312. <https://doi.org/10.1146/annurev-resource-100518-093954>
7. Martins KB, Faccioli PY, Bonesso MF, et al. Characteristics of resistance and virulence factors in different species of coagulase-negative staphylococci isolated from milk of healthy sheep and animals with subclinical mastitis. J Dairy Sci. 2017;100(3):2184–2195. <https://doi.org/10.3168/jds.2016-11583>
8. De Vlieghe S, Ohnstad I, Piepers S. Management and prevention of mastitis: A multifactorial approach with a focus on milking, bedding and data-management. J Integrative Agric. 2018;17(6):1214–1233. [https://doi.org/10.1016/S2095-3119\(17\)61893-8](https://doi.org/10.1016/S2095-3119(17)61893-8)
9. Leelahapongsathon K, Schukken YH, Srithanasuwan A, Suriyasathaporn W. Molecular epidemiology of *Streptococcus uberis* intramammary infections: Persistent and transient patterns of infection in a dairy herd. J Dairy Sci. 2020;103(4):3565–3576. <https://doi.org/10.3168/jds.2019-17281>
10. Huang C H, Furukawa K, Kusaba N, et al. Genetic parameters for novel mastitis traits defined by combining test-day somatic cell score and differential somatic cell count in the first lactation of Japanese Holsteins. J Dairy Sci. 2024; In Press, ISSN 0022-0302, <https://doi.org/10.3168/jds.2023-24399>
11. Damm M, Holm C, Blaabjerg M, et al. Differential somatic cell count. A novel method for routine mastitis screening in the frame of Dairy Herd Improvement testing programs. J Dairy Sci. 2017;100(6):4926–4940. <https://doi.org/10.3168/jds.2016-12409>
12. Rajčević M, Potočnik K, Levstek J. Correlations between somatic cells count and milk composition with regard to the season. Agric Consp Sci. 2003;68(3):221–226.
13. Nyman AK, Fasth C, Waller KP. Intramammary infections with different non-aureus staphylococci in dairy cows. J Dairy Sci. 2018;101(2):1403–1418. <https://doi.org/10.3168/jds.2017-13467>
14. De Visscher A, Piepers S, Haesebrouck F, et al. Teat apex colonization with coagulase-negative *Staphylococcus* species before parturition: distribution and species-specific risk factors. J Dairy Sci. 2016;99(2):1427–1439. <https://doi.org/10.3168/jds.2015-10326>
15. Duse A, Persson-Waller K, Pedersen K. Microbial aetiology, antibiotic susceptibility and pathogen-specific risk factors for udder pathogens from clinical mastitis in dairy cows. Animals. 2021;11(7):2113. <https://doi.org/10.3390/ani11072113>
16. Haw SR, Adkins PRF, Gosselin VB, et al. Intramammary infections in lactating Jersey cows: Prevalence of microbial organisms and association with milk somatic cell count and persistence of infection. J Dairy Sci. 2023; In Press, ISSN 0022-0302, <https://doi.org/10.3168/jds.2023-23848>
17. Meijering A, Hogeveen H, De Koning CJAM. Automatic milking, a better understanding: A Better Understanding. Wageningen Academic Publishers. 2004;544, ISBN: 978-90-76998-38-1. <https://doi.org/10.3920/978-90-8686-525-3>
18. Kaskous S. Physiological aspects of milk somatic cell count in dairy cattle. Int J Livest Res. 2021;1:1–12. eISSN: 2277-1964
19. European Commission. Commission Implementing Regulation (EU) No 853/2014 of 5 August 2014 repealing Regulation (EC) No 1151/2009 imposing special conditions governing the import of sunflower oil originating in or consigned from Ukraine. Off J Eur. Union 2014, L233, 25.
20. International Dairy Federation. Suggested interpretation of mastitis terminology. IDF Bulletin 338. Brussels, Belgium. 1999;3–26.
21. Miklós L, Trembos P, Hrnčiarova T. Limitations of intensive agricultural activity by types of abiotic complexes. 2002; 1: 1 500 000; 1: 1 500 000.
22. STN EN ISO 13366-2. Milk. Determination of the number of somatic cells. Part 2: Instructions for use of equipment for electronic particle counting using the fluorescence optical method (ISO 13366-2: 2006), Office for Standardization, Metrology and Testing of the Slovak Republic. 2007.
23. Li N, Richoux R, Boutinaud M, et al. Role of somatic cells on dairy processes and products: a review. Dairy sSci Technol. 2014;94:517–538. <https://doi.org/10.1007/s13594-014-0176-3>
24. Costa A, De Marchi M, Sagrafoli D, et al. Milk somatic cell count and polymorphonuclear cells in healthy quarters of cows that

- underwent blanket and selective dry therapy. Italian case study. *Vet Sci.* 2021;8(12):298. <https://doi.org/10.3390/vetsci8120298>
25. Godden SM, Royster E, Timmerman J, et al. Evaluation of an automated milk leukocyte differential test and the California Mastitis Test for detecting intramammary infection in early- and late-lactation quarters and cows. *J Dairy Sci.* 2017;100(8):6527–6544. <https://doi.org/10.3168/jds.2017-12548>
 26. Rudolf M, Scherer S. High incidence of *Listeria monocytogenes* in European red smear cheese. *Int J Food Microbiol.* 2001;63:91–98. [https://doi.org/10.1016/S0168-1605\(00\)00413-X](https://doi.org/10.1016/S0168-1605(00)00413-X)
 27. Jayarao BM, Pillai SR, Sawant AA. Guidelines for monitoring bulk tank milk somatic cell and bacterial counts. *J Dairy Sci.* 2004;87(10):3561–3573. [https://doi.org/10.3168/jds.S0022-0302\(04\)73493-1](https://doi.org/10.3168/jds.S0022-0302(04)73493-1)
 28. Věříš M. Neglected causal agents of mastitis. *Veterinářství,* 2019;69(9):592–594.
 29. Collier R. Dairy research: “Real” science or marketing? 2016 188 (10) 715–716; <https://doi.org/10.1503/cmaj.109-5278>
 30. Thompson JS, Green MJ, Hyde R, et al. The use of machine learning to predict somatic cell count status in dairy cows post-calving. *Front Vet Sci.* 2023;10. <https://doi.org/10.3389/fvets.2023.1297750>
 31. Pelmuş RŞ, Grosu H, Gras MA, et al. Estimation of the genetic parameters for Somatic Cell Scores in the first lactation of Romanian Black and White cattle. *Archiva Zootechnica.* 2022;25(1):142–153. <https://doi.org/10.2478/azibna-2022-0010>
 32. Whist AC, Osteras O. Associations between somatic cell counts at calving or prior to drying-off and clinical mastitis in the remaining or subsequent lactation. *J Dairy Res.* 2007;74:66–73. <https://doi.org/10.1017/S0022029906002172>
 33. Andersen S, Dohoo IR, Riekerink RO, et al. Diagnosing intramammary infections: Evaluating expert opinions on the definition of intramammary infection using conjoint analysis. *J Dairy Sci.* 2010;93(7):2966–2975. <https://doi.org/10.3168/jds.2009-2726>
 34. Samková E. Milk: production and quality: Milk: production and quality: scientific monograph, 1st ed. České Budějovice: University of South Bohemia in České Budějovice, Faculty of Agriculture, 2012. ISBN 978-80-7394-383-7
 35. Kvapilík J, Kučera J, Bucek P, et al. Yearbook of cattle breeding in the Czech Republic. MJP Online Early <http://admin.cmsch.cz/store/rocenka-chovu-skotu> (access: 2017.04.10)
 36. Riekerink RO, Barkema HW, Stryhn H. The effect of season on somatic cell count and the incidence of clinical mastitis. *J Dairy Sci.* 2007;90(4):1704–1715. <https://doi.org/10.3168/jds.2006-567>
 37. Wang Y, Nan X, Zhao Y, et al. Consumption of supplementary inulin modulates milk microbiota and metabolites in dairy cows with subclinical mastitis. *Appl Environ Microbiol.* 2022;88(4):e02059–21. <https://doi.org/10.1128/aem.02059-21>
 38. Sharma N, Singh NK, Bhadwal MS. Relationship of somatic cell count and mastitis: An overview. *Asian-Australasian J Animal Sci.* 2011;24(3):429–438.
 39. De Koning K, Slaghuis B, Van der Vorst Y. Robotic milking and milk quality: effects on bacterial counts, somatic cell counts, freezing point and free fatty acids. *Italian J Animal Sci.* 2003;2(4):291–299.
 40. Gygax L, Neuffer I, Kaufmann C, et al. Comparison of functional aspects in two automatic milking systems and auto-tandem milking parlors. *J Dairy Sci.* 2007;90(9):4265–4274. <https://doi.org/10.3168/jds.2007-0126>
 41. Schwarz D, Kleinhans S, Reimann G, et al. Investigation of dairy cow performance in different udder health groups defined based on a combination of somatic cell count and differential somatic cell count. *Prev Vet Med.* 2020;183(105123). <https://doi.org/10.1016/j.prevetmed.2020.105123>
 42. Huang CH, Furukawa K, Kusaba N. Estimating the nonlinear interaction between somatic cell score and differential somatic cell count on milk production by parity using generalized additive models. *J Dairy Sci.* 2023; 106(11):7942–7953. <https://doi.org/10.3168/jds.2022-22958>
 43. Zazharska N, Fotina T, Yatsenko I, Tarasenko L, Biben I, Zazharskyi V, et al. Comparative analysis of the criteria for goat milk assessment in Ukraine and France. *Ukr J Ecol.* 2021;11(2):144–148. https://doi.org/10.15421/2021_91
 44. Fotina T, Fotina H, Ladyka V, Ladyka L, Zazharska N. Monitoring research of somatic cells count in goat milk in the eastern region of Ukraine. *J Hellenic Vet Med Soc.* 2018;69(3):1101–1108. <https://doi.org/10.12681/jhvms.18882>